

Superconducting gaps in FeSe studied by soft point-contact Andreev reflection spectroscopy

Naidyuk Y., Kvitnitskaya O., Gamayunova N., Bashlakov D., Tyutrina L., Fuchs G., Hühne R., Chareev D., Vasiliev A.

Kazan Federal University, 420008, Kremlevskaya 18, Kazan, Russia

Abstract

© 2017 American Physical Society. FeSe single crystals have been studied by soft point-contact Andreev reflection spectroscopy. Superconducting gap features in the differential resistance $dV/dI(V)$ of point contacts such as a characteristic Andreev reflection double-minimum structure have been measured versus temperature and magnetic field. Analyzing dV/dI within the extended two-gap Blonder-Tinkham-Klapwijk model allows one to extract both the temperature and magnetic field dependence of the superconducting gaps. The temperature dependence of both gaps is close to the standard BCS behavior. Remarkably, the magnitude of the double-minimum structure gradually vanishes in magnetic field, while the minima position only slightly shifts with field, indicating a weak decrease of the superconducting gaps. Analyzing the $dV/dI(V)$ spectra for 25 point contacts results in the averaged gap values $(\Delta_L)=1.8\pm0.4\text{meV}$ and $(\Delta_S)=1.0\pm0.2\text{meV}$ and reduced values $2(\Delta_L)/kBT_c=4.2\pm0.9$ and $2(\Delta_S)/kBT_c=2.3\pm0.5$ for the large (L) and small (S) gap, respectively. Additionally, the small gap contribution was found to be within tens of percent, decreasing with both temperature and magnetic field. No signatures in the dV/dI spectra were observed, testifying to a gapless superconductivity or the presence of even smaller gaps.

<http://dx.doi.org/10.1103/PhysRevB.96.094517>

References

- [1] A. I. Coldea and M. D. Watson, The key ingredients of the electronic structure of FeSe, arXiv:1706.00338;
- [2] Yu. V. Pustovit and A. A. Kordyuk, Metamorphoses of electronic structure of FeSe-based superconductors (Review Article), *Fiz. Nizk. Temp.* 42, 1268 (2016)
- [3] Yu. V. Pustovit and A. A. Kordyuk, [*Sov. J. Low Temp. Phys.* 42, 995 (2016)]. 10.1063/1.4969896
- [4] S. Kasahara, T. Watashige, T. Hanaguri, Y. Kohsaka, T. Yamashita, Y. Shimoyama, Y. Mizukami, R. Endo, H. Ikeda, K. Aoyama, T. Terashima, S. Uji, T. Wolf, H. v. Löhneysen, T. Shibauchi, and Y. Matsuda, Field-induced superconducting phase of FeSe in the BCS-BEC crossover, *Proc. Natl. Acad. Sci. USA* 111, 16309 (2014). PNASA6 0027-8424 10.1073/pnas.1413477111
- [5] C. J. Chen, *Introduction to Scanning Tunneling Microscopy*, 2nd ed. (Oxford University Press, Oxford, UK, 2007);
- [6] J. E. Hoffman, Spectroscopic scanning tunneling microscopy insights into Fe-based superconductors, *Rep. Prog. Phys.* 74, 124513 (2011). RPPHAG 0034-4885 10.1088/0034-4885/74/12/124513
- [7] Yu. G. Naidyuk and I. K. Yanson, *Point-Contact Spectroscopy*, Springer Series in Solid-State Sciences Vol. 145 (Springer, New York, 2005).
- [8] D. Daghero and R. S. Gonnelli, Probing multiband superconductivity by point-contact spectroscopy, *Supercond. Sci. Technol.* 23, 043001 (2010). SUSTEF 0953-2048 10.1088/0953-2048/23/4/043001

- [9] C.-L. Song, Y.-L. Wang, P. Cheng, Y.-P. Jiang, W. Li, T. Zhang, Z. Li, K. He, L. Wang, J.-F. Jia, H.-H. Hung, C. Wu, X. Ma, X. Chen, and Q.-K. Xue, Direct observation of nodes and twofold symmetry in FeSe superconductor, *Science* 332, 1410 (2011). SCIEAS 0036-8075 10.1126/science.1202226
- [10] T. Watashige, Y. Tsutsumi, T. Hanaguri, Y. Kohsaka, S. Kasahara, A. Furusaki, M. Sigrist, C. Meingast, T. Wolf, H. V. Löhneysen, T. Shibauchi, and Y. Matsuda, Evidence for Time-Reversal Symmetry Breaking of the Superconducting State near Twin-Boundary Interfaces in FeSe Revealed by Scanning Tunneling Spectroscopy, *Phys. Rev. X* 5, 031022 (2015). 2160-3308 10.1103/PhysRevX.5.031022
- [11] S. A. Moore, J. L. Curtis, C. Di Giorgio, E. Lechner, M. Abdel-Hafiez, O. S. Volkova, A. N. Vasiliev, D. A. Chareev, G. Karapetrov, and M. Iavarone, Evolution of the superconducting properties in (Equation presented), *Phys. Rev. B* 92, 235113 (2015). PRBMDO 1098-0121 10.1103/PhysRevB.92.235113
- [12] L. Jiao, C.-L. Huang, S. Rößler, C. Koz, U. K. Rößler, U. Schwarz, and S. Wirth, Superconducting gap structure of FeSe, *Sci. Rep.* 7, 44024 (2017). 2045-2322 10.1038/srep44024
- [13] S. V. Borisenko, D. V. Evtushinsky, Z.-H. Liu, I. Morozov, R. Kappenberger, S. Wurmehl, B. Büchner, A. N. Yaresko, T. K. Kim, M. Hoesch, T. Wolf, and N. D. Zhigadlo, Direct observation of spin-orbit coupling in iron-based superconductors, *Nat. Phys.* 12, 311 (2016). 1745-2473 10.1038/nphys3594
- [14] J. Hong and D. S. L. Abergé, A universal explanation of tunneling conductance in exotic superconductors, *Sci. Rep.* 6, 31352 (2016). 10.1038/srep31352
- [15] Ya. G. Ponomarev, S. A. Kuzmichev, M. G. Mikheev, M. V. Sudakova, S. N. Tchesnokov, T. E. Shanygina, O. S. Volkova, A. N. Vasiliev, and Th. Wolf, Andreev spectroscopy of FeSe: Evidence for two gap superconductivity, *J. Exp. Theor. Phys.* 113, 459 (2011). JTPHES 1063-7761 10.1134/S1063776111080097
- [16] D. Chareev, E. Osadchii, T. Kuzmicheva, J.-Y. Lin, S. Kuzmichev, O. Volkova, and A. Vasiliev, Single crystal growth and characterization of tetragonal (Equation presented) superconductors, *CrystEngComm* 15, 1989 (2013). 1466-8033 10.1039/c2ce26857d
- [17] Yu. G. Naidyuk, N. V. Gamayunova, O. E. Kvitnitskaya, G. Fuchs, D. A. Chareev, and A. N. Vasiliev, Analysis of nonlinear conductivity of point contacts on the base of FeSe in the normal and superconducting state, *Fiz. Nizk. Temp.* 42, 42 (2016)
- [18] Yu. G. Naidyuk, N. V. Gamayunova, O. E. Kvitnitskaya, G. Fuchs, D. A. Chareev, and A. N. Vasiliev, [Sov. J. Low Temp. Phys. 42, 31 (2016)]. 10.1063/1.4939153
- [19] Yu. G. Naidyuk, G. Fuchs, D. A. Chareev, and A. N. Vasiliev, Doubling of the critical temperature of FeSe observed in point contacts, *Phys. Rev. B* 93, 144515 (2016). 2469-9950 10.1103/PhysRevB.93.144515
- [20] Y. J. Song, J. B. Hong, B. H. Min, Y. S. Kwon, K. J. Lee, M. H. Jung, and J.-S. Rhyee, Superconducting properties of a stoichiometric FeSe compound and two anomalous features in the normal state, *J. Korean Phys. Soc.* 59, 312 (2011). KPSJAS 0374-4884 10.3938/jkps.59.312
- [21] A. Bhaskar, H.-J. Huang, and C.-J. Liu, Effects of Mn doping on the normal-state transport of tetragonal FeSe superconductor up to 700 K, *Europhys. Lett.* 108, 17011 (2014). EULEEJ 0295-5075 10.1209/0295-5075/108/17011
- [22] G. Sheet, S. Mukhopadhyay, and P. Raychaudhuri, Role of critical current on the point-contact Andreev reflection spectra between a normal metal and a superconductor, *Phys. Rev. B* 69, 134507 (2004). 10.1103/PhysRevB.69.134507
- [23] A. A. Sinchenko, P. D. Grigoriev, A. P. Orlov, A. V. Frolov, A. Shakin, D. A. Chareev, O. S. Volkova, and A. N. Vasiliev, Gossamer high-temperature bulk superconductivity in FeSe, *Phys. Rev. B* 95, 165120 (2017). 2469-9950 10.1103/PhysRevB.95.165120
- [24] See Supplemental Material at <http://link.aps.org/supplemental/10.1103/PhysRevB.96.094517> for the comparison with literature data and for the statistical graphics.
- [25] Yu. G. Naidyuk, O. E. Kvitnitskaya, L. V. Tiutrina, I. K. Yanson, G. Behr, G. Fuchs, S.-L. Drechsler, K. Nenkov, and L. Schultz, Peculiarities of the superconducting gaps and the electron-boson interaction in (Equation presented) as seen by point-contact spectroscopy, *Phys. Rev. B* 84, 094516 (2011). PRBMDO 1098-0121 10.1103/PhysRevB.84.094516
- [26] P. O. Sprau, A. Kostin, A. Kreisel, A. E. Böhmer, V. Taufour, P. C. Canfield, S. Mukherjee, P. J. Hirschfeld, B. M. Andersen, and J. C. Séamus Davis, Discovery of orbital-selective Cooper pairing in FeSe, *Science* 357, 75 (2017). SCIEAS 0036-8075 10.1126/science.aal1575
- [27] B. I. Verkin, I. K. Yanson, I. O. Kulik, O. I. Shklyarevskii, A. A. Lysykh, and Yu. G. Naidyuk, Singularities in (Equation presented) dependences of point contacts between ferromagnetic metals, *Solid State Commun.* 30, 215 (1979). SSCOA4 0038-1098 10.1016/0038-1098(79)90337-5
- [28] Yu. G. Naidyuk, N. N. Gribov, O. I. Shklyarevskii, A. G. M. Jansen, and I. K. Yanson, Thermoelectric effects and the asymmetry of the current-voltage characteristic of metallic point contacts, *Fiz. Nizk. Temp.* 11, 1053 (1985)
- [29] Yu. G. Naidyuk, N. N. Gribov, O. I. Shklyarevskii, A. G. M. Jansen, and I. K. Yanson, [Sov. J. Low Temp. Phys. 11, 580 (1985)].

- [30] Yu. G. Naidyuk, O. E. Kvitnitskaya, I. K. Yanson, G. Fuchs, S. Haindl, M. Kidszun, L. Schultz, and B. Holzapfel, Point-contact study of (Equation presented)(Re=La, Sm) superconducting films, *Supercond. Sci. Technol.* 24, 065010 (2011). SUSTEF 0953-2048 10.1088/0953-2048/24/6/065010
- [31] Yu. G. Naidyuk, O. E. Kvitnitskaya, S. Aswartham, G. Fuchs, K. Nenkov, and S. Wurmehl, Exploring point-contact spectra of (Equation presented) in the normal and superconducting states, *Phys. Rev. B* 89, 104512 (2014). PRBMDO 1098-0121 10.1103/PhysRevB.89.104512
- [32] A. V. Chubukov, I. Eremin, and D. V. Efremov, Superconductivity versus bound-state formation in a two-band superconductor with small Fermi energy: Applications to Fe pnictides/chalcogenides and doped (Equation presented), *Phys. Rev. B* 93, 174516 (2016). 2469-9950 10.1103/PhysRevB.93.174516
- [33] J.-Y. Lin, Y. S. Hsieh, D. A. Chareev, A. N. Vasiliev, Y. Parsons, and H. D. Yang, Coexistence of isotropic and extended (Equation presented)-wave order parameters in FeSe as revealed by low-temperature specific heat, *Phys. Rev. B* 84, 220507 (R) (2011). PRBMDO 1098-0121 10.1103/PhysRevB.84.220507